

Number of probable and confirmed communicable disease cases in North Carolina by disease for: 1) the current month, 2) the average cases for the year to date, 3) the total cases during 2011, 4) and the average (with 95% confidence intervals) of the previous five years (2006 to 2010).

DISEASE	Cases in January, 2012	Average Cases in January, 2007 - 2011	Cases in 2011	Average cases (95% confidence interval) per year 2006 to 2010
Botulism ¹	0	0	2	1 (0 – 4)
Campylobacter Infection*	62	36	909	693 (361 – 1,026)
Chlamydia ²	5,723	2,396	54,894 	39,161 (29,695 – 48,641)
Cryptosporidiosis	0	2	115	114 (22 – 206)
E. coli O157:H7/ other STEC Infection*	5	8	155	126 (57 – 194)
Ehrlichiosis ³	1	1	107	67 (40 – 174)
Gonorrhea	1,791	1,092	17,487	15,336 (8,779 – 21,892)
Group A Strep Infection, Invasive	24	15	181	143 (81 – 205)
Haemophilus Influenzae	9	7	84	86 (7 – 179)
Hepatitis A	3	4	31	60 (11 – 109)
Hepatitis B (acute)	5	8	124	128 (66 – 190)
Hepatitis B (perinatal)	0	0	1	2 (0 – 5)
Hepatitis B (chronic) ⁴	75	76	1,309 	873 (662 – 1,085)
Hepatitis C (acute)	3	2	61	29 (6 – 64)
Influenza Death, Adult ⁵	0	3	26	50 (0 – 152)
Influenza Death, Pediatric	0	1	10	3 (0 – 14)
LaCrosse Encephalitis	0	0	24	15 (1 – 31)
Legionellosis	4	3	86 	52 (15 – 85)
Listeriosis	1	1	21	25 (8 – 42)
Lyme Disease	1	2	75	66 (23 – 155)

DISEASE	Cases in January, 2012	Average Cases in January, 2007 - 2011	Cases in 2011	Average cases (95% confidence interval) per year 2006 to 2010
Malaria ⁶	1	3	49	33 (2 – 68)
Measles	0	0	2	1 (0 – 4)
Meningococcal Invasive Disease	0	2	15	22 (4 – 48)
Mumps	0	0	9	20 (0 – 88)
Pelvic Inflammatory Disease	59	44	677 	445 (214 - 647)
Pertussis	4	21	185	287 (121 – 453)
Spotted Fever Rickettsiosis	3	6	332	496 (131 – 1,124)
Rubella	0	0	1	0 (0)
Salmonellosis*	106	80	2,517	1,862 (1,019 – 2,705)
Shigellosis*	4	14	225	237 (89 – 563)
Toxic Shock Syndrome ⁷	0	0	16	10 (0 – 26)
Typhoid Fever	0	1	8	7 (0 – 14)
Vibrio Infections	0	1	15	19 (1 – 38)
West Nile Encephalitis	0	0	2	2 (0 – 12)

Does not include HIV, Syphilis and TB. ¹Infant, foodborne and wound botulism cases combined; ²Chlamydia annual case average calculated for 2008-2010; ³Includes HE, HME and unspecified; ⁴Represents an artificial increase in 2011 due to review and disposition of 2008-2010 cases; ⁵Influenza-associated adult deaths became reportable in 2010. ⁶All cases are imported; ⁷Includes non-streptococcal and streptococcal infections. *Per CSTE case definition, includes suspect cases.

 = significant increase (≥ 3 standard deviations above average).  = significant decrease (≥ 3 standard deviations below average).

Reportable communicable diseases with NO reported cases in the current year were not included in this report. Because cases are routinely updated, case numbers may change (data was extracted on 2/21/12). Case definitions for these diseases are available at: <http://epi.publichealth.nc.gov/cd/lhds/manuals/cd/toc.html>.



North Carolina Division of Public Health
 Epidemiology Section, Communicable Disease Branch
 Megan Davies, State Epidemiologist

