

Hepatitis B, Chronic

2012 Case Definition

CSTE Position Statement Number: 11-ID-04

Clinical Description

No symptoms are required. Persons with chronic hepatitis B virus (HBV) infection may have no evidence of liver disease or may have a spectrum of disease ranging from chronic hepatitis to cirrhosis or liver cancer.

Laboratory Criteria for Diagnosis

- Immunoglobulin M (IgM) antibodies to hepatitis B core antigen (IgM anti-HBc) negative AND a positive result on one of the following tests: hepatitis B surface antigen (HBsAg), hepatitis B e antigen (HBeAg), or nucleic acid test for hepatitis B virus DNA (including qualitative, quantitative and genotype testing), OR
- HBsAg positive or nucleic acid test for HBV DNA positive (including qualitative, quantitative and genotype testing) or HBeAg positive two times at least 6 months apart (Any combination of these tests performed 6 months apart is acceptable)

Case Classification

Probable

A person with a single HBsAg positive or HBV DNA positive (including qualitative, quantitative and genotype testing) or HBeAg positive lab result and does not meet the case definition for acute hepatitis B.

Confirmed

A person who meets either of the above laboratory criteria for diagnosis.

Comment

Multiple laboratory tests indicative of chronic HBV infection may be performed simultaneously on the same patient specimen as part of a "hepatitis panel." Testing performed in this manner may lead to seemingly discordant results, e.g., HBsAg-negative AND HBV DNA-positive. For the purposes of this case definition, any positive result among the three laboratory tests mentioned above is acceptable, regardless of other testing results. Negative HBeAg results and HBV DNA levels below positive cutoff level do not confirm the absence of HBV infection.